

LISTING OF CLAIMS

1. (Currently amended) A method for accelerating the cell cycle of a cell, comprising delivering to a cell an effective amount of electromagnetic energy to accelerate the cell cycle of said cell at least 2 fold, wherein said cell is part of a gastrointestinal tissue, ~~synovium~~, or lung tissue.
2. (Original) The method of claim 1, wherein the rate at which said cell replicates its DNA increases.
3. (Original) The method of claim 1, wherein the G1 stage of said cell cycle is shortened.
4. (Canceled)
5. (Original) The method of claim 1, wherein said electromagnetic energy has a wavelength in a region of the spectrum selected from the group consisting of X-ray radiation, ultraviolet radiation, visible radiation, infrared radiation, microwave radiation and radiofrequency radiation.
6. (Original) The method of claim 1, wherein said electromagnetic energy comprises an energy that is in the range of 1 to 300 mW/cm².
7. (Original) The method of claim 1, wherein said electromagnetic energy is pulsed.
8. (Previously presented) The method of claim 1, wherein said cell is selected from the group consisting of fibroblast, neuronal cell, epithelial cell, macrophage, neutrophil, keratinocyte, endothelial cell, epidermal melanocyte, hair follicle papilla cell, skeletal muscle

cell, smooth muscle cell, osteoblast, neuron, chondrocyte, hepatocyte, pancreatic cell, kidney cell, aortic cell, bronchial cell and tracheal cell.

9. (Currently amended) The method of claim 1, further comprising delivering to said cell an effective amount of electromagnetic energy to activate a cell cycle regulator.

10. (Currently amended) The method of claim 1, ~~further comprising delivering to~~ wherein said ~~cell~~ an effective amount of electromagnetic energy is sufficient to activate a signal transduction protein.

11. (Currently amended) The method of claim 1, ~~further comprising delivering to~~ wherein said ~~cell~~ an effective amount of electromagnetic energy is sufficient to activate a transcription factor.

12. (Currently amended) The method of claim 1, ~~further comprising delivering to~~ wherein said ~~cell~~ an effective amount of electromagnetic energy is sufficient to activate a DNA synthesis protein.

13. (Currently amended) The method of claim 1, ~~further comprising delivering to~~ wherein said ~~cell~~ an effective amount of electromagnetic energy is sufficient to activate a receptor.

14. (Currently amended) The method of claim 1, ~~further comprising delivering to~~ wherein said ~~cell~~ an effective amount of electromagnetic energy is sufficient to inhibit an Angiotensin Receptor.

15. (Currently amended) A method for activating a cell cycle regulator, comprising delivering to a cell an effective amount of electromagnetic energy to activate said cell cycle

regulator, wherein said cell cycle regulator accelerates the cell cycle of said cell, and wherein said cell is part of a gastrointestinal tissue, ~~synovium~~, or lung tissue.

16. (Canceled)

17. (Previously presented) The method of claim 15, wherein the rate at which said cell replicates its DNA increases.

18. (Previously presented) The method of claim 15, wherein the G1 stage of said cell cycle is shortened.

19. (Previously presented) The method of claim 15, wherein said cell cycle is accelerated 2 fold.

20. (Original) The method of claim 15, wherein said electromagnetic energy has a wavelength in a region of the spectrum selected from the group consisting of X-ray radiation, ultraviolet radiation, visible radiation, infrared radiation, microwave radiation and radiofrequency radiation.

21. (Original) The method of claim 15, wherein said electromagnetic energy comprises an energy that is in the range of 1 to 300 mW/cm².

22. (Original) The method of claim 15, wherein said electromagnetic energy is pulsed.

23. (Original) The method of claim 15, wherein said cell is selected from the group consisting of a fibroblast, neuronal cell, epithelial cell, macrophage, neutrophil, keratinocyte, endothelial cell, epidermal melanocyte, hair follicle papilla cell, skeletal muscle cell, smooth

muscle cell, osteoblast, neuron, chondrocyte, hepatocyte, pancreatic cell, kidney cell, aortic cell, bronchial cell and tracheal cell.

24. (Currently amended) A method for activating a signal transduction protein, comprising delivering to a cell an effective amount of electromagnetic energy to activate said signal transduction protein, and wherein said cell is part of a gastrointestinal tissue, ~~synovium~~, or lung tissue.

25. (Currently amended) A method for activating a transcription factor, comprising delivering to a cell an effective amount of electromagnetic energy to activate said transcription factor, and wherein said cell is part of a gastrointestinal tissue, ~~synovium~~, or lung tissue.

26. (Currently amended) A method for activating a DNA synthesis protein, comprising delivering to a cell an effective amount of electromagnetic energy to activate said DNA synthesis protein, and wherein said cell is part of a gastrointestinal tissue, ~~synovium~~, or lung tissue.

27. (Currently amended) A method for activating a receptor, comprising delivering to a cell an effective amount of electromagnetic energy to activate said receptor, and wherein said cell is part of a gastrointestinal tissue, ~~synovium~~, or lung tissue.

28. (Currently amended) A method for inhibiting an angiotensin receptor, comprising delivering to a cell an effective amount of electromagnetic energy to inhibit said angiotensin receptor, and wherein said cell is part of a gastrointestinal tissue, ~~synovium~~, or lung tissue.

29. - 62. (Canceled)

63. (Currently amended) A method for accelerating the cell cycle, comprising delivering to a cell an effective amount of electromagnetic energy to accelerate the cell cycle of said cell at least 10 percent, and wherein said cell is part of a gastrointestinal tissue, ~~synovium~~, or lung tissue.

64. (Currently amended) A method for accelerating the cell cycle, comprising delivering to a cell an effective amount of electromagnetic energy to accelerate the cell cycle of said cell at least 25 percent, and wherein said cell is part of a gastrointestinal tissue, ~~synovium~~, or lung tissue.

65. (Previously presented) The method of claim 64, wherein the rate at which said cell replicates its DNA increases.

66. (Previously presented) The method of claim 64, wherein the G1 stage of said cell cycle is shortened.

67. (Previously presented) The method of claim 64, wherein said electromagnetic energy has a wavelength in a region of the spectrum selected from the group consisting of X-ray radiation, ultraviolet radiation, visible radiation, infrared radiation, microwave radiation and radiofrequency radiation.

68. (Previously presented) The method of claim 64, wherein said electromagnetic energy comprises an energy that is in the range of 1 to 300 mW/cm².

69. (Previously presented) The method of claim 64, wherein said electromagnetic energy is pulsed.

70. (Previously presented) The method of claim 64, wherein said cell is selected from the group consisting of fibroblast, neuronal cell, epithelial cell, macrophage, neutrophil, keratinocyte, endothelial cell, epidermal melanocyte, hair follicle papilla cell, skeletal muscle cell, smooth muscle cell, osteoblast, neuron, chondrocyte, hepatocyte, pancreatic cell, kidney cell, aortic cell, bronchial cell and tracheal cell.

71. (Currently amended) The method of claim 64, ~~further comprising delivering to wherein~~ said cell an effective amount of electromagnetic energy is sufficient to activate a cell cycle regulator.

72. (Currently amended) The method of claim 64, ~~further comprising delivering to wherein~~ said cell an effective amount of electromagnetic energy is sufficient to activate a signal transduction protein.

73. (Currently amended) The method of claim 64, ~~further comprising delivering to wherein~~ said cell an effective amount of electromagnetic energy is sufficient to activate a transcription factor.

74. (Currently amended) The method of claim 64, ~~further comprising delivering to wherein~~ said cell an effective amount of electromagnetic energy is sufficient to activate a DNA synthesis protein.

75. (Currently amended) The method of claim 64, ~~further comprising delivering to wherein~~ said cell an effective amount of electromagnetic energy is sufficient to activate a receptor.

76. (Currently amended) The method of claim 64, ~~further comprising delivering to~~ wherein said cell an effective amount of electromagnetic energy is sufficient to inhibit an Angiotensin Receptor.

77.-84. (Canceled).

85. (Currently amended) A method for accelerating the cell cycle, comprising delivering to a cell an effective amount of electromagnetic energy to accelerate the cell cycle of said cell at least 50 percent, and wherein said cell is part of a gastrointestinal tissue, ~~synovium~~, or lung tissue.

86. (Currently amended) A method for accelerating the cell cycle, comprising delivering to a cell an effective amount of electromagnetic energy to accelerate the cell cycle of said cell at least 75 percent, and wherein said cell is part of a gastrointestinal tissue, ~~synovium~~, or lung tissue.